

تحية شكر وتقدير الي الزملاء اطباء (ادارة مديرية الطب البيطري) محافظة القاهره







تحية شكر وتقدير الي:
الأخت الزميله الكريمه العزيزه (الدكتوره ناديه وديع)





تحية شكر وتقدير الي: الدكتور السيد عبيد (وكيل الوزاره ومدير الطب البيطري محافظة القاهره)


علاجات الطفيليات في الكلاب والقطط



أهم طفيليات الكلاب




أهم طفيليات القطط



الطفيليات الداخليه

INTESTINAL PARASITES IN DOGS



أهم الادويه المستعمله للطفيليات الداخليه



أهم الأدويه المستعمله للطفيليات الخارجيه




مستقبل أدويه الكلاب والقطط




الطفيليات الداخليه

INTESTINAL PARASITES IN DOGS



- Internal Parasites
 - Roundworms
 - Live in dogs intestines
 - Consume partially digested feed
 - Passed by feces
 - Offspring to offspring







Symptoms of Roundworms

- There will be no symptoms if there are just a few worms present
- If there are hundreds, the puppy will become "potbellied"
- Worms will live on the puppies nutrients.
- Puppies should be treated for worms at 2, 4, 6 and 8 weeks.
- Should treat all puppies like they have worms.





Medscape® www.medscape.com



Source: Infect Med © 2003 Cliggott Publishing, Division of SCP Communications

- Internal Parasites
 - Hookworms
 - Small worm like parasites that hooks on the intestinal wall
 - Feeds on blood of host
 - Contract orally, through skin, pass from off spring to off spring



- Hook Worm "Symptoms"
 - Pale Gums
 - Diarrhea
 - Weakness
 - Failure to grow properly
- Hook Worm "Treatment"
 - Drugs (Oral or Injection)
 - Drugs only kill adult hookworm
 - One adult will lay 20, 000 larvae
 - Human can not contract



- Roundworms "Symptoms"
 - Loss of appetite
 - Pot Belly
 - Vomiting
 - Diarrhea
- Roundworms "Treatment"
 - Dewormer (2 4 Weeks)





أهم الادويه المستعمله للطفيليات الداخليه





Plantain martin per silverment products between the



Panacul[®] (fenbendazole)

Dewormer

Granules 22.2% (222 mg/g)

FOR DOGS AND FOR ONLY THE CARNIVOROUS/OMNIVOROUS ANIMALS LISTED ON THE SIDE PANEL

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

1 lb (454 g)

Retain carton for complete product information.



RETREATMENT:

Schedule may vary depending on the climate where you live and the activity of your dog.

The following schedule should be used as a general guideline.

Weaned pups (6-8 weeks of age) should be dewormed at 6, 8, 10 and 12 weeks of age. Treat the dam at the same time as the puppies. Dogs over 6 months of age should be dewormed at

SIDE EFFECTS:

FENCUR is safe for use in puppies and kittens 6 weeks or older and adults, including pregnant. Do not deworm a puppy that is sick. Consult your veterinarian for diagnosis of the illness. In U.S. clinical studies 3 of 240 dogs had vomiting associated with the use of the product.



DOGS

41-50 lbs 2 1/2 capsules 51-60 lbs 3 capsules 61-80 lbs 4 capsules

DIRECTIONS:

Give capsule whole or mix with a small amount of wet food. Make sure your dog eats all the medicated food. You may need to moisten dry dog food to aid mixing. REPEAT DOSE 3 DAYS IN A ROW.

Marine Company Control of the William

FDA APPROVED EXP. DATE 3/20 QUEST LABORATORIES P.O. BOX 1815 Corpus Christi TX 78480

DOSAGE CHART


DOG WEIGHT	PACKET SIZE
10 LBS	1 gram
20 LBS	2 gram
30 LBS*	1 gram + 2 gram
40 LBS	4 gram
50 LBS*	1 gram + 4 gram
60 LBS*	2 gram + 4 gram
80 LBS*	4 gram + 4 gram
80+ LBS	* Use combination pouches to obtain recommended daily dose

SAFE FOR ALL DOGS

- House Pets
- » Pregnant Dogs
- Show Dogs
- > Heartworm-Infected Dogs
- > Elderly Dogs
- Young Puppies over the age of 6 weeks > Working Dogs




"Pathology" Active Segments on Pet




"Pathology" Active Segments on Poop







Cysticercoid Larvae



<u>Pathology</u>

- Definitive Host (Dog & Cat)
 - No Pathology (nutrient competition in mal-nourished hosts)
 - "Client Worry" (proglottid aesthetics)
- Intermediate Host (Flea)
 - (Who cares?)

Clinical Signs

- Segments presented by Client
- Pet usually shows no signs
- Occasionally pet drags tail
- Observation of fleas
- History
 - Lack of Flea Control
 - Access to stray dogs / cats

Diagnosis

- Segment Squash
 - Oval segment
 - Egg Packets





أهم الادويه المستعمله للطفيليات الداخليه









Treatment

- Praziquantel (Droncit) (5mg/kg) [dog & cat]
- Pyrantel+Praziquantel+Febantel (Drontal Plus Tablets) [dog]
- Pyrantel+Praziquantel (Virbantel Flavored Chewables) [dog] & (Drontal Tablets) [cat]
- Ivermectin+Pyrantel+Praziquantel (Iverhart Max Chewable Tablets) [dog]
- Emodepside+Praziquantel (Profender) [cat]
- Epsiprantel (Cestex) [dog & cat]

الطفيليات الخارجيه



- External Parasites
- External Parasites -
 - Live outside the host
 - Skin
 - Coat





What do ticks look like?



- External Parasites
 - Tick
 - Small parasites that feeds on host
 - Will bury itself into skin
 - May carry many other disease (lymes)
 - Feed off host and fall off





- Ticks "Symptoms"
 - Ticks physically present
 - Lymes disease paralysis, arthritis
- Tick "Treatment"
 - Physical removal
 - Tick shampoo
 - Lymes -No treatment, but drugs to relieve symptoms.



أهم الأدويه المستعمله للطفيليات الخارجيه







For dogs 5.1-10 lbs in weight and at least 6 weeks of age

Six single-dose tubes each containing 0.25 mL (30 mg) of selamectin



Administer once a month to prevent heartworm disease and prevent and control flea infestations

OPEN

HERE



Caution: U.S. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

For use in animals only. Keep out of reach of children. NADA WH52, Approved by FDA 6 Doses




- Flea "Symptoms"
 - Scratching and biting
 - Red spots and rashes
- Flea "Treatment"
 - Flea shampoo
 - Medication (internal and external)
 - Extermination



- External Parasites
 - Fleas
 - Small fly/nat looking parasite
 - Will lay egg in coat
 - Bit animal to feed on blood









مستقبل أدويه الكلاب والقطط







الزملاء الكرام بمديرية طب بيطري القاهره



شكرا لحضوركم الكريم

thank



